

Bromhexine acts by:

- A. Inhibiting cough centre
- B. Irritating gastric mucosa and reflexly increasing bronchial secretion
- C. Depolymerizing mucopolysaccharides present in sputum.
- D. Desensitizing stretch receptors in the lungs

Dextromethorphan is an:

- A. Analgesic
- B. Antitussive
- C. Expectorant
- D. Antihistaminic

The following antitussive is present in opium but has no analgesic or addicting properties:

- A. Noscapine
- B. Codeine
- C. Pholcodeine
- D. Ethylmorphine

Budesonide is a:

- A. Nonsteroidal antiinflammatory drug
- B. High ceiling diuretic
- C. Inhaled corticosteroid for asthma
- D. Contraceptive

One of the most common side effect of inhaled beclomethasone dipropionate is:

- A. Pneumonia
- B. Oropharyngeal candidiasis
- C. Atrophic rhinitis
- D. Pituitary-adrenal suppression

Which of the following has overdose toxicity that includes insomnia, arrhythmias, and convulsions?

- (a) Aminophylline
- (b) Cromolyn
- (c) Epinephrine
- (d) Ipratropium

Which of the following is a very long acting β_2 – selective agonist that is used for asthma prophylaxis?

- (a) Aminophylline
- (b) Cromolyn
- (c) Salmeterol
- (d) Ipratropium

Antitussives act by:

- (a) Liquifying bronchial secretions
- (b) Raising the threshold of cough centre
- (c) Reducing cough inducing impulses from the lungs
- (d) Both (b) and (c)

The following is **not a first line antitubercular drug**

- a) Isoniazid
- b) Ethambutol
- c) Rifampin
- d) Cycloserine

Pyridoxine is given prophylactically with the following drug to prevent neurotoxicity

- a) Isoniazid
- b) Ethambutol
- c) Streptomycin
- d) Pyrazinamide

The following drug causes orange red discolouration of urine and secretions:

- a) Isoniazid
- c) Ethambutol
- b) Rifampin
- d) Cycloserine

The following drug is related to sulfonamides and acts by inhibition of folate synthase.

- a) Ethionamide
- b) Kanamycin
- c) Cycloserine
- d) Para-aminosalicylic acid

In multy drug resistant TB The following drug is added:

- | | |
|---------------|-----------------|
| a) Kanamycin | b) Amikacin |
| c)Capreomycin | d)Ciprofloxacin |

The following drug is used in Chemoprophylaxis of TB

- | | |
|---------------|-----------------------------|
| a)Cycloserine | b) Para-aminosalicylic acid |
| c)Isoniazid | d) Amikacin |